



			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		yesadme
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		yesadme
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		

1346
1364
1379
1398
1472
1576
1579
1623
1645
2650
4691
5007
5208
5320
5345
5441
5641

N. Froio Degori. BEHAVIORAL AND ELECTROCORTICAL CHANGES INDUCED BY PARAQUAT AFTER INJECTION IN SPECIFIC AREAS OF THE BRAIN OF THE RAT. <i>Neuropharmacology</i> . 1988. 27:201-207
Jayshree P Nellore. Paraquat exposure induces behavioral deficits in larval zebrafish during the window of dopamine neurogenesis. <i>Toxicology Reports</i> . 2015. 2:950-956
Q. Chen, Y. Niu, R. Zhang, H. Guo, Y. Gao, Y. Li, R. Liu. The toxic influence of paraquat on hippocampus of mice: involvement of oxidative stress. <i>Neurotoxicology</i> . 2010. 31:310-6
J. H. Ellwanger, P. Molz, D. R. Dallemole, A. Pereira dos Santos, T. E. Muller, L. Cappelletti, M. Goncalves da Silva, S. I. Franke, D. Pra, J. A. Pegas Henriques. Selenium reduces bradykinesia and DNA damage in a rat model of Parkinson's disease. <i>Nutrition</i> . 2015. 31:359-65
K. M. Costa, I. S. Maciel, L. W. Kist, M. M. Campos, M. R. Bogo. Pharmacological inhibition of CXCR2 chemokine receptors modulates paraquat-induced intoxication in rats. <i>PLoS One</i> . 2014. 9:e105740
L. Miranda-Contreras, R. Davila-Ovalles, P. Benitez-Diaz, Z. Pena-Contreras, E. Palacios-Pru. Effects of prenatal paraquat and mancozeb exposure on amino acid synaptic transmission in developing mouse cerebellar cortex. <i>Brain Res Dev Brain Res</i> . 2005. 160:19-27
Y. H. Sun, Y. Li, Y. J. Niu, Q. Chen, R. Zhang. [Effects of paraquat on the learning and memory ability in developing mice]. <i>Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi</i> . 2011. 29:437-9
D. E. Woolley, D. W. Gietzen, S. J. Gee, J. Magdalou, B. D. Hammock. Does paraquat (PQ) mimic MPP+ toxicity?. <i>Proc West Pharmacol Soc</i> . 1989. 32:191-3
M. E. Nunes, T. E. Muller, M. M. Braga, B. D. Fontana, V. A. Quadros, A. Marins, C. Rodrigues, C. Menezes, D. B. Rosemberg, V. L. Loro. Chronic Treatment with Paraquat Induces Brain Injury, Changes in Antioxidant Defenses System, and Modulates Behavioral Functions in Zebrafish. <i>Mol Neurobiol</i> . 2016. #volume#: #pages#
H. F. Li, S. X. Zhao, B. P. Xing, M. L. Sun. Ulinastatin suppresses endoplasmic reticulum stress and apoptosis in the hippocampus of rats with acute paraquat poisoning. <i>Neural Regen Res</i> . 2015. 10:467-72
Y. Pham Pesah. Drosophila parkin mutants have decreased mass and cell size and increased sensitivity to oxygen radical stress. <i>Development</i> . 2004. 131:2183-94
Y. Ait-Bali, S. Ba-M'hamed, M. Bennis. Prenatal Paraquat exposure induces neurobehavioral and cognitive changes in mice offspring. <i>Environ Toxicol Pharmacol</i> . 2016. 48:53-62
A. Kumar, P. K. Christian, K. Panchal, B. R. Guruprasad, A. K. Tiwari. Supplementation of Spirulina (Arthrospira platensis) Improves Lifespan and Locomotor Activity in Paraquat-Sensitive DJ-1betaDelta93 Flies, a Parkinson's Disease Model in Drosophila melanogaster. <i>J Diet Suppl</i> . 2017. 14:573-588
D. Quintero-Espinosa, M. Jimenez-Del-Rio, C. Velez-Pardo. Knockdown transgenic Lrrk Drosophila resists paraquat-induced locomotor impairment and neurodegeneration: A therapeutic strategy for Parkinson's disease. <i>Brain Res</i> . 2017. 1657:253-261
R. M. Satpute, P. P. Pawar, S. Puttewar, S. D. Sawale, P. D. Ambhore. Effect of resveratrol and tetracycline on the subacute paraquat toxicity in mice. <i>Hum Exp Toxicol</i> . 2017. #volume#:960327116688070
S. K. Yadav, S. N. Rai, S. P. Singh. Mucuna pruriens reduces inducible nitric oxide synthase expression in Parkinsonian mice model. <i>J Chem Neuroanat</i> . 2017. 80:1-10
L. J. Yoo S. E. Na R. Liu Y. H. Ran Q. T. Chen. Cognitive impairment and increased A beta levels induced by paraquat exposure are attenuated by enhanced removal of mitochondrial H2O2. <i>Neurobiology of Aging</i> . 2012. 33: #pages#

Exposure to environmental risk factors such as herbicides in early life has been proposed to play important roles in the d
Environmental paraquat (PQ) exposure has been suggested to be a potential risk factor for neurodegenerative disorders
OBJECTIVE: The aim of this study was to explore the effects of selenium (Se) on locomotor activity and DNA damage in a
Paraquat (PQ) is an agrochemical agent commonly used worldwide, which is allied to potential risks of intoxication. This
The goal of this study was to analyze the effects of prenatal exposure to the pesticides paraquat (PQ) and mancozeb (MZ
OBJECTIVE: To explore the damages of paraquat to the learning and memory ability of developing mice and explore the p
Paraquat (PQ) administration consists in a chemical model that mimics phenotypes observed in Parkinson's disease (PD),
Lung injury is the main manifestation of paraquat poisoning. Few studies have addressed brain damage after paraquat po
Mutations in the gene parkin in humans (PARK2) are responsible for a large number of familial cases of autosomal-recess
In the present work, we investigated developmental toxicity of Paraquat (PQ), from the 1st or 6th day of mating and thro
Spirulina (Arthrospira platensis) is a cyanobacterium (blue-green alga) consumed by humans and other animals because
Leucine-rich repeat kinase 2 (LRRK2) has been linked to familial and sporadic Parkinson's disease. However, it is still unre
Paraquat (PQ) is a nonselective bipyridyl herbicide widely used in agriculture to control weeds, but its accidental, occupa
Parkinson's disease is one of the most common neurodegenerative disease found in aged peoples. Plentiful studies are b
Pesticide exposure is a risk factor of Alzheimer's disease (AD). However, little is known about how pesticide exposure ma

Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant



			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		



5739
5842
6218
6230
6426
15
27
36
58
95
106
108
113
120
133

M. Ninkovic M. Stevanovic I. Ilic K. Durdevic D. Dukic. The effect of pre-treatment with l-name on glutathione and glutathione peroxidase in paraquat-induced neurotoxicity in rat striatum. <i>Arhiv za Farmaciju</i> . 2012. 62:237-251
E. Chronis N. Gourgou. Chemically induced oxidative stress affects ASH neuronal function and behavior in <i>C. elegans</i> . <i>Scientific Reports</i> . 2016. 6:#pages#
T. E. Nunes M. E. Menezes C. C. Marins A. T. Leitemperger J. Gressler A. C. L. Carvalho F. B. de Freitas C. M. Quadros V. A. Fachinetto R. Rosemberg D. B. Loro V. L. Müller. Sodium Selenite Prevents Paraquat-Induced Neurotoxicity in Zebrafish. <i>Molecular Neurobiology</i> . 2017. #volume#:1-14
J. Shajan A. Antony D. Cynthia Pauline P. Karthick Raja Namasivayam S. Nellore. Proteinaceous compounds from fragaria ananassa fruit attenuates paraquat induced Parkinson like locomotor and mitochondrial alterations in Zebrafish. <i>International Journal of Pharmacy and Pharmaceutical Sciences</i> . 2015. 7:246-251
J. Sirapat W. Thiantanawat A. Satayavivad. Neurological effects of chronic exposure to low doses of paraquat in rats. <i>Research Communications in Pharmacology and Toxicology</i> . 1997. 2:269-282
V. N. Li J. Fink A. L. Uversky. Pesticides directly accelerate the rate of alpha-synuclein fibril formation: a possible factor in Parkinson. <i>FEBS Lett</i> . 2001. 500:105-8
B. K. Lee D. W. Cory-Slechta D. A. Opanashuk L. A. Barlow. Modulation of antioxidant defense systems by the environmental pesticide maneb in dopaminergic cells. <i>Neurotoxicology</i> . 2004. 26:63-75
C. Ko H. S. Thomas B. Tsang F. Chew K. C. Tay S. P. Ho M. W. Lim T. M. Soong T. W. Pletnikova O. Troncoso J. Dawson V. L. Dawson T. M. Lim K. L. Wang. Stress-induced alterations in parkin solubility promote parkin aggregation and compromise parkin's protective function. <i>Hum Mol Genet</i> . 2005. 14:3885-97
K. Y. Korlipara L. V. Cooper J. M. Schapira A. H. Chau. Protection against paraquat and A53T alpha-synuclein toxicity by cabergoline is partially mediated by dopamine receptors. <i>J Neurol Sci</i> . 2008. 278:44-53
M. Ba M. Liang H. Ma L. Yu Q. Yu T. Wang Y. Kong. 5'-Aza-dC sensitizes paraquat toxic effects on PC12 cell. <i>Neurosci Lett</i> . 2012. 524:35-9
J. R. Obando D. Liu J. Ganio G. Volitakis I. Mok S. S. Crouch P. J. White A. R. Codd R. Liddell. Lipophilic adamantyl- or deferrioxamine B have enhanced neuroprotective capacity: implications for Parkinson disease. <i>Free Radic Biol Med</i> . 2013. 60:147-56
A. Betemps D. Morignat E. Lazizzera C. Hogeveen K. Andrieu T. Baron T. Chorfa. Specific pesticide-dependent increases in alpha-synuclein levels in human neuroblastoma (SH-SY5Y) and melanoma (SK-MEL-2) cell lines. <i>Toxicol Sci</i> . 2013. 133:289-97
S. K. Tewari D. Tetrud J. W. Langston J. W. Schule B. Mak. Mitochondrial dysfunction in skin fibroblasts from a Parkinson's disease patient with an alpha-synuclein triplication. <i>J Parkinsons Dis</i> . 2011. 1:175-83
S. D. Dolatabadi N. Chan S. F. Zhang X. Akhtar M. W. Parker J. Soldner F. Sunico C. R. Nagar S. Talantova M. Lee B. Lopez K. Nutter A. Shan B. Molokanova E. Zhang Y. Han X. Nakamura T. Masliah E. Yates J. R. rd Nakanishi N. Andreyev A. Y. Okamoto S. Jaenisch R. Ambasudhan R. Lipton S. A. Ryan. Isogenic human iPSC Parkinson's model shows nitrosative stress-induced dysfunction in MEF2-PGC1alpha transcription. <i>Cell</i> . 2013. 155:1351-64
Y. Erkut C. Pan-Montojo F. Boland S. Stewart M. P. Muller D. J. Wurst W. Hyman A. A. Kurzchalia T. V. Toyoda. Products of the Parkinson's disease-related glyoxalase DJ-1, D-lactate and glycolate, support mitochondrial membrane potential and neuronal survival. <i>Biol Open</i> . 2014. 3:777-84

Contact herbicide paraquat (DQ) is bipyridylum compound, which undergoes redox metabolism; hence enlarged in hum
Oxidative stress (OS) impact on a single neuron's function in vivo remains obscure. Using <i>C. elegans</i> as a model organism
Considering the antioxidant properties of sodium selenite ( $\text{Na}_2\text{SeO}_3$ ) and the involvement of oxidative stress events in p
Objectives: To assess the Parkinson like locomotor and mitochondrial alterations, associated with the exposure of paraqu
The neurotoxic effects of chronic exposure to subcutaneous administration of low doses of paraquat on motor behavior a
Parkinson's disease involves intracellular deposits of alpha-synuclein in the form of Lewy bodies and Lewy neurites. The e
A lack of evidence supporting a role of heritability in the development of idiopathic Parkinson's disease (PD) has implicat
Mutations in parkin are currently recognized as the most common cause of familial Parkinsonism. Emerging evidence als
Both genetic and environmental factors are thought to be involved in the aetiology of Parkinson's disease (PD). Oxidative
Parkinson's disease (PD) is one of the common neurodegenerative diseases that result in the progressive damage of dop
Parkinson disease (PD) is a neurodegenerative disease characterized by death of dopaminergic neurons in the substantia
Epidemiological studies indicate a role of genetic and environmental factors in Parkinson's disease involving alterations c
Mitochondrial dysfunction has been frequently implicated in the neurodegenerative process that underlies Parkinson's d
Parkinson's disease (PD) is characterized by loss of A9 dopaminergic (DA) neurons in the substantia nigra pars compacta (
Parkinson's disease is associated with mitochondrial decline in dopaminergic neurons of the substantia nigra. One of the

Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant



			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
			Animal	aniprim	anisecc		
						invitro	
						invitro	
						invitro	
						invitro	
						invitro	
						invitro	
						invitro	
						invitro	
						invitro	

141
170
207
322
323
334
337
341
343
350
351
357
358
374
378
379
381
383
402
405
410
419

F. F. Carretta D. Lattanzio F. Palmisano M. Candeletti S. Romualdi P. Caputi. Proteasome subunit and opioid receptor gene expression down-regulation induced by paraquat and maneb in human neuroblastoma SH-SY5Y cells. <i>Environ Toxicol Pharmacol.</i> 2015. 40:895-900
X. F. Li S. Chou A. P. Bronstein J. M. Wang. Inhibitory effects of pesticides on proteasome activity: Implication in Parkinson's disease. <i>Neurobiology of Disease.</i> 2006. 23:198-205
E. Lascala Janda. Parkinsonian toxin-induced oxidative stress inhibits basal autophagy in astrocytes via NQO2/quinone oxidoreductase 2: Implications for neuroprotection. <i>Autophagy.</i> 2015. 11:1063-80
G. Schluter Schmuck. An in vitro model for toxicological investigations of environmental neurotoxins in primary neuronal cell cultures. <i>Toxicol Ind Health.</i> 1996. 12:683-96
D. Miyako Kang. Changes of energy metabolism induced by 1-methyl-4-phenylpyridinium (MPP+)-related compounds in rat pheochromocytoma PC12 cells. <i>Arch Biochem Biophys.</i> 1997. 337:75-80
M. Bauer Vogt. Oxidative stress and hypoxia/reoxygenation trigger CD95 (APO-1/Fas) ligand expression in microglial cells. <i>FEBS Lett.</i> 1998. 429:67-72
W. L. Sun Yang. Paraquat-induced cell death in PC12 cells. <i>Neurochem Res.</i> 1998. 23:1387-94
K. Tokunaga Morita. Cytotoxic effect of paraquat on rat C6 glioma cells: evidence for the possibility of non-oxidative damage to the cells. <i>Jpn J Pharmacol.</i> 1999. 79:121-4
X. Sun Li. Paraquat induced activation of transcription factor AP-1 and apoptosis in PC12 cells. <i>J Neural Transm (Vienna).</i> 1999. 106:1-21
H. Possel Noack. Peroxynitrite mediated damage and lowered superoxide tolerance in primary cortical glial cultures after induction of the inducible isoform of NOS. <i>Glia.</i> 1999. 28:13-24
B. J. Patel Day. A mechanism of paraquat toxicity involving nitric oxide synthase. <i>Proc Natl Acad Sci U S A.</i> 1999. 96:12760-5
S. I. Pakbaz Said. NMDA receptor activation: critical role in oxidant tissue injury. <i>Free Radic Biol Med.</i> 2000. 28:1300-2
G. Ahr Schmuck. Rat cortical neuron cultures: an in vitro model for differentiating mechanisms of chemically induced neurotoxicity. <i>In Vitro Mol Toxicol.</i> 2000. 13:37-50
H. S. Gibson Chun. Dopaminergic cell death induced by MPP(+), oxidant and specific neurotoxicants shares the common molecular mechanism. <i>J Neurochem.</i> 2001. 76:1010-21
Q. Keller Ding. Proteasome inhibition in oxidative stress neurotoxicity: implications for heat shock proteins. <i>J Neurochem.</i> 2001. 77:1010-7
L. Marchese Rossi. Increased susceptibility of copper-deficient neuroblastoma cells to oxidative stress-mediated apoptosis. <i>Free Radic Biol Med.</i> 2001. 30:1177-87
M. R. Aquilano Ciriolo. Differential role of superoxide and glutathione in S-nitrosoglutathione-mediated apoptosis: a rationale for mild forms of familial amyotrophic lateral sclerosis associated with less active Cu,Zn superoxide dismutase mutants. <i>J Neurochem.</i> 2001. 77:1433-43
Q. Y. Pedersen Li. Dependence of excitotoxic neurodegeneration on mitochondrial aconitase inactivation. <i>J Neurochem.</i> 2001. 78:746-55
F. Hashino Osakada. Neuroprotective effects of alpha-tocopherol on oxidative stress in rat striatal cultures. <i>Eur J Pharmacol.</i> 2003. 465:15-22
N. P. Politi Rotstein. Protective effect of docosahexaenoic acid on oxidative stress-induced apoptosis of retina photoreceptors. <i>Invest Ophthalmol Vis Sci.</i> 2003. 44:2252-9
G. Rohrdanz Schmuck. Oxidative stress in rat cortical neurons and astrocytes induced by paraquat in vitro. <i>Neurotox Res.</i> 2002. 4:1-13
A. Stolzing Ernst. Antioxidants effectively prevent oxidation-induced protein damage in OLN 93 cells. <i>Arch Biochem Biophys.</i> 2004. 421:54-60



Paraquat (PQ) and maneb (MB) are able to induce neurotoxic effects by promoting alpha-synuclein (alpha-syn) aggregate
Epidemiological studies have suggested a correlation of pesticides and Parkinson's disease (PD) while genetic and bioche
Oxidative stress (OS) stimulates autophagy in different cellular systems, but it remains controversial if this rule can be ge
Currently, most neurotoxicological investigations are still conducted using various animal models (e.g. chickens, rodents)
We examined effects of three structurally related pyridinium compounds, 1-methyl-4-phenylpyridinium (MPP+), paraqua
Apoptosis plays an important role in neurodegeneration, although the mechanisms and mediators in the brain are largel
Paraquat was taken up by PC12 cells in a carrier-mediated, saturable manner. When PC12 cells were permeabilized with
Although paraquat has been shown to cause oxidative damage to neuronal cells, little is known about its effect on glial ce
Drugs and certain environmental toxins may be responsible for the pathogenesis of Parkinson's disease. We have used p
The effect of the induction of i-NOS in primary glial cultures was studied with respect to the protein levels of reactive oxy
Paraquat (PQ) is a well described pneumotoxicant that produces toxicity by redox cycling with cellular diaphorases, there
The excitatory amino acid glutamate serves important neurologic functions, but overactivation of its N-methyl-D-asparta
Various structurally unrelated chemicals [2,5 hexandione, acrylamide, organophosphates like mipafox, beta,beta iminodi
Recent etiological study in twins (Tanner et al. 1999) strongly suggests that environmental factors play an important role
Recent studies have demonstrated that inhibition of the proteasome, an enzyme responsible for the majority of intracell
Treatment of neuroblastoma cells with the copper chelator triethylene tetramine tetrahydrochloride induced intracellular
SH-SY5Y cells transfected with the enzymatically inactive Cu,Zn superoxide dismutase mutant H46R were more resistant
Using the inactivation of mitochondrial and cytosolic aconitases as markers of compartment-specific superoxide (O <sub>2</sub> <sup>-</sup> ) p
Oxidative stress caused by an increase in free radicals plays an important role in neuronal death. We investigated the eff
PURPOSE: In a recent study, it was demonstrated that docosahexaenoic acid (DHA) promotes the survival of retinal phot
Oxidative stress has been discussed as crucial mechanism of neuronal cell death in the adult brain. However, it was not c
Oxidative stress is supposed to play an important role in demyelinating diseases. Oligodendrocytes are the myelin-formi

Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant
Relevant